

STIC-ILL

From: Arthur, Lisa
Sent: Monday, November 05, 2001 2:22 PM
To: STIC-ILL
Subject: reference for 08/484,786

370403
QR184.I5
NOS

4907310

Please provide a copy of the following:

1. Conner et al. (1983) PNAS 80: 278-282.
2. Rollini et al. PNAS (1985 Nov) 82(12): 7197-7201.
3. Gorski et al. IMMUNOGENTICS (1987) 25(6):379-402.
4. de Preval et al. IMMUNOGEENTICS (1987) 26(4-5): 249-257
5. Irle et al. J. EXPERIMENTAL MEDICINE (1988 Mar 1) 167(3): 853-872.
6. Andersson et al. IMMUNOGENETICS (1988) 28 (1): 1-5.

Thanks,
Lisa Arthur
CM1 12E13
1655
308-3988

Scientific and Technical
Information Center

NOV 06 RECD 4/02

PAT. & T.M. OFFICE

CONFIDENTIAL

2820873

DNA typing of *HLA-DR_β* chain genes can discriminate between undetected alleles and real homozygotes

C. de Préal¹*, G. Angelini^{1, 2}, B. Boogh¹, G.B. Ferrara², and B. Mach¹

¹ Department of Microbiology, School of Medicine, University of Geneva, C.M.U., 9, Avenue de Champel, CH-1211 Geneva, Switzerland

² Istituto Nazionale per la Ricerca sul Cancro, Genova, Italy and Centro Ricerche Immunoematologiche A.V.I.S., Bergamo, Italy.

Abstract. The polymorphism of HLA-DR antigens has been studied by Southern blot hybridization under conditions specific for the detection of the *DR_β* chain genes. Haplotype-specific patterns were defined with DNA from *DR1*, 2, 3, 4, 7, *w8*, *w11*, *w12*, and *W13* homozygous typing cells, with restriction enzymes *Eco* RI, *Bgl* I, and *Pvu* II. Certain serological specificities, such as *DR2*, *DR3*, and *DR7*, can be encoded by distinct allelic forms of *DR_β* chain genes. The procedure of "DNA typing" was applied to family analysis of individuals expressing only a single DR specificity upon serological typing. Three cases are described here: (1) in family GR, phenotypic *DR 7* homozygotes correspond to genomic heterozygotes, and a novel *DR7* allele is described; (2) in family RU, the genes corresponding to a serologically undetected (blank) *DR* allele were identified by restriction fragment length polymorphism (RFLP); this novel *DR* haplotype has an RFLP pattern similar to those of the *DRw52* family, even though this specificity was not expressed on the DR-blank lymphocytes; (3) in family RG, there is no blank allele, but a homozygote RFLP situation at the *DR* subregion.

Introduction

The *HLA-D* region of the human major histocompatibility complex (MHC) codes for class II or Ia antigens. These class II antigens are the products of MHC-linked immune response genes which control the ability to trigger an immune response against foreign antigens (Nagy et al. 1981). The Ia molecules consist of two transmembrane glycoproteins, the α (M_r 33 000–35 000) and the β (M_r 27 000–29 000) chains. In man three biochemically dis-

tinct groups of polymorphic class II antigens have been described, HLA-DR, -DQ, and -DP, encoded in three distinct subregions within the MHC (Kaufman et al. 1984, Mach et al. 1986). An important characteristic of Ia molecules is the extensive allelic polymorphism which is responsible for the self-restriction of cell-cell interaction within immunocompetent cells of each individual (Schwartz 1985). This polymorphism is of importance for organ transplantation since matching of class II loci, and of HLA-DR in particular, is essential for graft survival (Opelz 1984). Another relevant aspect of this polymorphism is that certain haplotypes are linked to susceptibility to certain diseases (reviewed by Möller 1983).

The established methodology for the analysis of HLA class II polymorphism involves serological, immunochemical, and cellular reagents (see reviews by Albert et al. 1984). All these procedures rely on the detection of the various epitopes of class II antigens and characterize the polymorphism and the complexity of the MHC system at the phenotypic level. The methodology of gene cloning and sequencing has led to a more direct analysis of the polymorphism and complexity of class II genes at the genotypic level, based on hybridization of cloned HLA class II genes to DNA restriction fragments (Wake et al. 1982). This procedure relies on polymorphic differences at certain restriction-enzyme sites and on allelic differences in the length of DNA fragments specific for individual class II genes (RFLP). This technique of "DNA typing" was first used to detect polymorphism of *DR* and *DQ β* chain genes (Wake et al. 1982).

Each of the three subregions, *DR*, *DQ*, *DP*, consists of a number of loci encoding α and β chains (Spielman et al. 1984, Auffray and Strominger 1986, Mach et al. 1986). In the case of HLA-DR (Fig. 1), there is a single, nonpolymorphic, α chain gene and several polymorphic β chain genes (Long et al. 1983). These have been linked on a molecular map (Spies et al. 1985, Rollini et al. 1985), and they show more than 90% sequence homology.

The present study is focused on *HLA-DR_β* chain

* Present address: Unité de Recherches d'Immunologie d'Immunogénétique Humaine, INSERM U 100, CHU, Purpan, 31052 Toulouse Cedex, France




Fig. 1. Organization of *HLA-D* region and molecular map of the *DR β* gene fragment used as hybridization probe. The top line represents a map of the *HLA-D* region. The number of *DR β* chain genes is derived from a study by Rollini and co-workers (1985) and may vary in different haplotypes (Böhme et al. 1985). The middle line represents the human genomic fragment subcloned into pBR 322 and used as *DR β* probe in Southern experiments (Fig. 2). This human gene fragment contains the exons (filled boxes) coding for the first and the second extracellular domains and for the transmembrane region of the *DR β* polypeptide chain. A map with five enzymes is presented; the scale is in kilobases (kb).

genes and the experimental conditions were chosen to detect *DR* and not *DQ* or *DP* β chain genes. DNA patterns of homozygous typing cells (HTCs) derived from genetically homozygous donors (consanguineous marriages) were used to interpret RFLP patterns of heterozygous individuals in family studies. An interesting correlation was observed between certain restriction DNA patterns and the known serological supertypic specificities (*DRw52/DRw53*). This analysis was focused on the interpretation of family pedigrees in cases where one or several individuals express only a single *DR* serological specificity. Three observations were made: some individuals who are phenotypically homozygous were found to be genetically heterozygous with a "split" of the serological specificity at the DNA level; some individuals

with a "blank" haplotype by serology have indeed a serologically unrecognized *HLA-DR* allele with a novel characteristic RFLP pattern corresponding to the "blank" haplotype; finally, some phenotypically homozygous individuals are homozygous by DNA genotyping. These "homozygous" *HLA-DR* phenotypes can now be interpreted in light of the RFLP pattern and new undetected alleles can be identified at the DNA level.

Materials and methods

Source of DNA. The origin and immunological typing of cell lines and blood samples used are described in Table 1.

Table 1. *HLA* phenotype of cells used in hybridizations

Donors	Form	Name	Consang.	HLA						Origin	Source
				A	B	C	D	DR	DQ	DP	
HTC	MLV	+	w32	27	w2	w1	1	w1	w2	Leiden	Giphart
HTC	PHS	+	3	7	w7	w2	2	w1	w4	Leiden	Giphart
HTC	AVL	+	1	8	w7	w3	3, w52	w2	w4	Leiden	Giphart
HTC	QBL	+	26	18	w5	w3	3, w52	w2	ws2	Leiden	Giphart
HTC	HAR	+	1	8	w7	w3	3, w52	w2	w4	Leiden	Giphart
HTC	CAA	+	1	8	na	w3	3, w52	w2	w2	Leiden	Giphart
HTC	BSM	+	2	w62	w3	w4	4, w53	w3	w2	Leiden	Giphart
HTC	ATH	+	25	18	na	w11	5, w52	w3	w2	Leiden	Giphart
HTC	HHK	+	3	7	w7	w18	w13, w52	w1	w4	Leiden	Giphart
HTC	MANN	+	29	44	na	w7	7, w53	w2	w2	Copenhagen	Svegaard
HTC	BM9	+	2	w35	w4	w8	w8	na	na	Bergamo	Ferrara
HTC	BM16	+	2	18	w7	DB6	w12, w52	na	na	Bergamo	Ferrara
LCL	H9	-	1, 2	7, 8	na	w2, 3	2, 3	w1, 2	w1, 2	Bethesda	Shaw
LCL	M14	-	1, 2	7, 8	na	w2, 3	2, 4	w1, 2	w1, 2	Bethesda	Shaw
LCL	PM1	-	1, 2	7, 8	na	w2, 3	2, 3	w1, 2	w1, 2	Bethesda	Shaw
LCL	R5	-	1, 2	7, 8	na	w2, 3	2, 3	w1, 2	w1, 2	Bethesda	Shaw
F	RG	see legend to Fig. 7								Bergamo	Ferrara
F	RU	see legend to Fig. 6								Bergamo	Ferrara
F	GR	see legend to Fig. 5								Bergamo	Ferrara

HTC, homozygous typing cell; LCL, lymphoblastoid cell line (EBV) (transformed); F, donors from a family; Consang., donor issued from consanguineous marriage (first cousins); na, not available

HLA-DR

Human gene
of Kunkel

DNA digest
pletion with
lambda DN
with either
11 are gene

Probe used
structed by
derived from
al. 1985). A
1. This hug
of the *DR β*
extracellular

Electrophor
depurinated
method of
and Schuel
Plus memb
used in tran

Prehybridiz
were rinsed
citrate). The
GeneScreen
were presoak
with gentle
1 x Denhard
acetate, 50 m
0.1% sodium
sperm DNA
buffer as use
translated D
twice for 30
SDS, 1 x De
0.1% SDS;
graphy at
about 3 day

Results and

Specificity
To achieve
DR haplot
covering a
control fo
Southern
genomic ge
was hybrid
BamG (Fig
the probe h
no cross-hy
Overloading
resulted in
(Fig. 2). I
signal is ab
der the cot
DR β probe

Human genomic DNA. DNA preparations were adapted from the method of Kunkel and co-workers (1977).

DNA digestions. Routinely 10 µg of genomic DNA was digested to completion with either Eco RI, Bgl I, Pvu II, or Pst I (6 units/µg⁻¹). Phage lambda DNA (0.6 µg/20 µl) were digested with 2 units/µg⁻¹ at 37 °C with either Eco RI or Hind III. The phages βDR 22, βDQ 18, and βDP 11 are genomic clones containing entire β chain genes.

Probe used for Southern hybridization. Plasmid pP14 BamG was constructed by subcloning into pBR322 a 4.3 kb Bam HI genomic fragment derived from a *DRw6β* gene cloned into phage Charon 30 (Gorski et al. 1985). A restriction map of the Bam HI insert is presented in Figure 1. This human DNA fragment contains almost all the coding sequence of the *DRβ* gene, including exons coding for the first and the second extracellular domains and for the transmembrane segment.

Electrophoresis and transfer of DNA fragments. DNA fragments were dephosphorylated, denatured, neutralized, and transferred according to the method of Southern (1975). Nitrocellulose filters (0.22 µ, Schleicher and Schuell, Dassel, Federal Republic of Germany) and GeneScreen Plus membrane (New England Nuclear, Boston, Massachusetts) were used in transfer experiments.

Prehybridization, hybridization, and washes. After transfer, the filters were rinsed with 4 × SSC (1 × SSC = 150 mM NaCl, 15 mM trisodium citrate). The DNA was fixed to nitrocellulose at 80 °C for 2 h, and to GeneScreen Plus by exposure to UV light (325 nm) for 2 min. Filters were presoaked in 5 × SSC, 5 × Denhardt's solution for 1 h at 68 °C with gentle shaking, and then prehybridized for 4–6 h at 68 °C in 1 × Denhardt's solution, 0.75 M NaCl, 5 mM ethylenediaminetetraacetate, 50 mM sodium phosphate buffer, pH 7.0, 10% dextran sulfate, 0.1% sodium dodecyl sulfate (SDS), 250 µg/ml⁻¹ denatured herring sperm DNA. Hybridization was carried out for 15 h at 68 °C in the same buffer as used for prehybridization with 1 × 10⁶ cpm/ml⁻¹ of ³²P nick-translated DNA probe (1–3 × 10⁸ cpm/µg⁻¹). Filters were washed twice for 30 min at 68 °C with the following solutions: 4 × SSC, 0.1% SDS, 1 × Denhardt's solution, 0.1% sodium pyrophosphate; 2 × SSC, 0.1% SDS; 0.2 × SSC, 0.1% SDS; 0.1 × SSC, 0.1% SDS. Autoradiography at –70 °C with a Cawo VI intensifying screen proceeded for about 3 days.

Results and discussion

Specificity of the detection of HLA-DR gene fragments.

To achieve maximum discrimination between different DR haplotypes, we made use of a large genomic probe covering almost the entire *DRβ* chain gene (Fig. 1). To control for the specificity of *DRβ* hybridization, a Southern blot was made using *DRβ*, *DQβ*, or *DPβ* genomic genes cloned in three different phages. The filter was hybridized with the nick-translated plasmid pP14 BamG (Fig. 2). Under the experimental conditions used, the probe hybridizes only with the cloned *DRβ* gene, and no cross-hybridization is visible with *DQ* or *DP* β genes. Overloading the gel with the same DR, DQ, and DP clones resulted in a slight cross-hybridization with the *DPβ* gene (Fig. 2). Densitometric analysis indicates that the *DPβ* signal is about 100-fold weaker than the *DRβ* signal. Under the conditions used to analyze genomic DNA, the *DRβ* probe hybridizes only to DR genes. In other studies,




Fig. 2. Specificity of hybridization with the *DRβ* chain probe pP14BamG. A Southern blot was made using cloned genes coding for *DRβ*, *DQβ*, and *DPβ* chain genes digested with Eco RI or Hind III (see Materials and methods). The filter was hybridized with the nick-translated *DRβ* chain plasmid pP14BamG (Fig. 1)

cross-hybridization of *DRβ* probes with *DQ* and *DPβ* chain genes has resulted in complex patterns which are often difficult to interpret at the molecular level (Cohen et al. 1984, Nicklas et al. 1985, Bell et al. 1985). It should be noted that the genomic probe pP14 BamG does not contain repetitive sequences and hybridizes only with *DRβ* gene fragments.

Definition and limits of allele-specific DNA patterns.

DNA from genetically homozygous cell lines (Table 1) was used as references to define allele-specific DNA patterns. The results obtained with Bgl I have already been published (Angelini et al. 1984), and Figure 3 represents a summary of the reference patterns obtained with the enzymes Bgl I, Eco RI, and Pvu II. Homologous restriction fragment patterns are presented side by side (Fig. 3). It is striking that groups of very similar DNA patterns can easily be recognized with each of the three enzymes used: DR 3, 5, w6, w12, and DR4, 7 respectively (Fig. 3). These two groups correspond to the well-known DRw52 and DRw53 supertypic groups characterized serologically (Tanigaki and Tosi 1982, Angelini et al. 1984). The haplotypes which could be classified in one of these two groups are therefore structurally related and they probably have a common origin in evolution (Gorski et al. 1986, 1987a). This observation may have some practical implications for an "intelligent" HLA-DR matching in organ transplanta-




Fig. 3. Diagrammatic patterns of DR β DNA fragment length of the homozygous typing cell lines (HTC) used as references. The origin and the typing of the DNA samples are summarized in Table 1 (AVL is used as DR 3/3 HTC). From left to right the DNA patterns were obtained with Bgl I, Pvu II, and Eco RI. These diagrams are based on Southern blot experiments as presented by Angelini and co-workers (1984). The *dark strips* represent major bands and the *dashed strips* represent fainter bands. The scale is in kilobases. A close similarity in two groups of DNA patterns is visible: DR 4/4, and DR 7/7 on one side (DRw53); DR3/3, DR5/5, DRw6/w6, and DRw12/w12 on the other side (DRw52)




Fig. 4. Examples of genomic heterogeneity within the same serological specificities. *Upper panel:* DNA in lanes 1, 2, 3, and 4 were prepared from DR3/3 genetically homozygous cell lines AVL, QBL, HAR, and CAA (Table 1). DNA samples were digested with Eco RI, Pvu II, Bgl I, and Bam HI. The circles indicate the position of DNA bands that are missing in sample 2 (QBL). *Lower panel:* Comparison of DR β chain gene restriction fragments from DR2/3 heterozygous individuals: Samples a, b, c, and d correspond to the cell lines H9, M14, PM1, and R5 (Table 1). The circles indicate the position of DR2-specific DNA bands that are missing in sample d. Reference homozygous patterns are included for comparison

tion. Ind
match ad
consequ
typic far
similar

In a
differed
Figure 3
to distin
of such
document
1984), a
at the D
DR2 and
an exam
paragrap
DR3 hos
lyzed wi
These fo
and Pvu
HI, sam
(AVL, H
map of
shows th
in QBL
lini et al
have sin
DNA see
preparat
gelini et




Fig. 5. Genomic heterogeneity of *DR_β* chain genes coding for the DR7 specificity. Haplotypes of family GR are: Father: A25, Bw51, Cw1, DR5, DRw52, DQw3. A29, Bw44, DR7, DRw53, DQw2. Mother: A2, Bw51, DR7, -, DQw3. A24, Bw44, Cw2, DR1, DQw1. DNA of the family members and of HTC DR1/1 and DR5/5 was digested with Bgl I. The DNA pattern transmitted by the DR7 haplotype of the father corresponds to the pattern of the HTC MANN (Fig. 3) and is represented as 7/7. The DNA pattern transmitted by the DR7 haplotype of the mother is different and is represented on the scheme as 7'/7'.

tion. Indeed we have proposed that a single HLA-DR mismatch across different supertypic groups will have greater consequences than a single mismatch within a given supertypic family, where the *DR* genes are structurally very similar (Mach et al. 1986).

In a few cases, we observed DNA patterns that differed slightly from the reference RFLP shown in Figure 3. Thus some serological specificities correspond to distinct alleles at the genomic level. The existence of such "splits" of serological specificities is well documented at the cellular level (Grosse-Wilde et al. 1984), at the biochemical level (Nepom et al. 1983), and at the DNA level (Se et al. 1985). Examples concerning *DR2* and *DR3* haplotypes are presented in Figure 4, and an example concerning *DR7* will be discussed in the next paragraph. DNA samples of four individuals known to be *DR3* homozygous (consanguineous marriages) were analyzed with four restriction enzymes (Fig. 4, upper panel). These four samples share the same distribution of Eco RI and Pvu II restriction sites. However, with Bgl I and Bam HI, sample 2 (QBL) differs from samples 1, 3, and 4 (AVL, HAR, CAA, Table 1). A study of the restriction map of the cloned *DR_β* chain genes of AVL (sample 1) shows that the 3.9 kb Bam HI fragment which is missing in QBL (sample 2) corresponds to the *DR_βIII* locus (Rolini et al. 1985). Indeed, the haplotypes of AVL and QBL have since been found to differ at the *DR_βIII* locus by DNA sequence analysis (J. Gorski et al., manuscript in preparation) and by oligonucleotide hybridization (Angelini et al. 1986a, Gorski et al. 1987b) corresponding

respectively to the *52a* and the *52b* allele at that locus (Gorski and Mach 1986). Another case of split of serological specificity with RFLP was observed on a *DR2* pattern. Four *DR2/3* DNA samples provided by S. Shaw were analyzed with Eco RI and Bgl I (Fig. 4, lower panel). In samples a, b, and c, DNA fragments correspond exactly to the sum of *DR2/2* and *DR3/3* reference patterns, whereas for sample d (cell line R5, Table 1), one *DR2*-specific fragment is missing on both Eco RI and Bgl I patterns. This difference may be correlated with a functional difference. Although cell line R5 is indistinguishable by serology from the other *DR2/3* cell lines H9, M14, and PM1 (Table 1) (Shaw et al. 1980), it produces a strong mixed lymphocyte reaction when tested with cells from the other *DR2/3* individuals (S. Shaw, personal communication). The split in *DR2* genotypes observed by RFLP (Fig. 4) correlates therefore with the functional differences observed in cellular reactivity. Additional polymorphism detected by RFLP is not necessarily relevant, however, since many polymorphic restriction sites are located in noncoding regions (introns and flanking regions). As discussed elsewhere (Mach et al. 1986), the analysis of polymorphic splits by hybridization with allele-specific oligonucleotides (Angelini et al. 1986a) does provide a way to identify phenotypically relevant micropolymorphism, down to the level of single nucleotide differences.

Discrimination between split of serological specificity, unknown *DR* allele, and homozygous *DR* genes. In many instances, only a single *DR* specificity is identified pheno-

n and the
lined with
The dark
A patterns

ume sero-
2, 3, and
cell lines
les were
ie circles
n sample
e restric-
Samples
'M1, and
'-specific
ozygous




Fig. 7. Analysis of family RG and DRw52, DQw2, the a/d haplotypes between haplotypes fragments from a haplotype

typically up if those individuals not, to identify show here studies. The In family G DR7 for one Three siblings

Fig. 6A and B of family RU. The DNA patterns Pvu II and in siblings). On the The numbers of the DR bands Pvu II and Bgl shared by DR




Fig. 7. Analysis at the DNA level of a DR-blank serological haplotype: example of a homozygous DR specificity. The serological haplotypes of family RG are: Father: *a*: A2, Bw44, Cw5, Bw4, DR5, DRw52. *b*: ? Mother: *c*: Aw24, B7, Bw6, DR2, DQw1. *d*: A2, Bw56, Cw1, Bw6, DR5, DRw52, DQw3. The serological typings indicate that three siblings, male and female, share the *a/c* haplotypes and two siblings, male and female, the *a/d* haplotypes. A single example of restriction patterns obtained with DNA from *a/c* or *a/d* offsprings is shown because no variation was found between haplotypically identical siblings. From left to right the DNA patterns were obtained with Eco RI, Bgl I, and Pst I. Open arrows correspond to DNA fragments from the DR5 haplotype and closed arrows to DNA fragments from the DR2 haplotype. The dot indicates that a given band can be used as a haplotype-specific marker in the context of the restriction fragments observed

typically upon HLA typing. It is essential to determine if those individuals are genotypically homozygous and if not, to identify the serologically undetected allele. We show here that this can be achieved by RFLP in family studies. Three different examples will be presented. (a) In family GR, the father and the mother were both typed DR7 for one of their haplotypes (see legend to Fig. 5). Three siblings inherited both DR7 alleles and were expect-

ed to be homozygous. However, as shown with Bgl I restriction fragments (Fig. 5), the parents differ in the structure of their DR7_β genes. The DR pattern of the father is identical with the DR7 of the cell line MANN, but the DR7 pattern of the mother lacks an 11 kb band and has an extra 4.7 kb fragment. In consequence, the DR7 haplotypes of the father and the mother are different at the DNA level (see 7/7 versus 7'/7' in Fig. 5) and the DR7/7 sib-

Fig. 6A and B. Analysis at the DNA level of DR-blank serological haplotype: Example of an undetected DR specificity. The serological haplotypes of family RU are: Father: A28, B47, DR5, DRw52, DQw3. A29, B44, DR7, DQw2. Mother: A2, B18, DQw3. A2, B5, Bw4, DRw8, DRw52j. The DNA patterns of family members and of three HTC used as references are presented in two panels: in A DNA digestion was performed with Pvu II and in B digestion was performed with Bgl I. The DR serological typing is indicated on the top of each lane (F, father; M, mother; s, siblings). On the right part of the figure the DNA patterns of all individuals with a DR-blank phenotype (5/-, 7/-, 8/-) are represented schematically. The numbers indicate the bands specific for DR 5, 7, or 8. The stars represent the additional DNA bands corresponding to the β chain genes of the DR blank haplotype. At the extreme right of the scheme the deduced DR DNA pattern of the blank haplotype (-/-) is represented both for Pvu II and Bgl I; the size of these genomic fragments is indicated on the left of the figure. On A it is not possible to know if the 1.9 kb fragment shared by DR5, DR7, and DR8 haplotypes is also present on the DR blank haplotype (?)

lings are therefore heterozygous (7/7'). The genomic pattern of the *DR7'* allele described here in the mother is also different from an additional *DR7* allele recently observed by RFLP by Angelini and co-workers (1986b) and therefore corresponds to a third *DR7* allele, characterized by a unique Bgl I pattern. Interestingly, the *DR7* haplotype of the mother (*DR7'*) does not encode the supertypic specificity generally associated with *DR7*, *DRw53*. In this haplotype, it is likely that the product of the locus encoding *DRw53* is not expressed, as suggested by the protein analysis of Karr (1986) in another *DR7*, *DRw53*⁻ situation. A deletion of the locus encoding *DRw53* has been suggested by the RFLP pattern in a similar case (Böhme et al. 1985).

(b) In family RU, serological analysis detected only a single *DR* allele in the mother and three siblings (see legend to Fig. 6). In these individuals, a DNA restriction fragment pattern corresponding to a novel, serologically "blank" haplotype could be identified. With the enzymes Pvu II and Bgl I, the DNA pattern of the *DR* blank haplotype was deduced from a comparison with the DNA of the control reference HTC's (Fig. 6). This comparison involved *DR5/5* with 5/- (sibling), *DR7/7* with 7/- (sibling), and *DR8/8* with 8/- (mother). This newly defined pattern of a *DR* blank haplotype consists of Pvu II bands at 13, 6.1, and 3.8 kb, and of Bgl I bands at 8.8, 7.8, and 2.4 kb. Interestingly, some of these bands are also found in the RFLP pattern of haplotypes belonging to the *DRw52* supertypic group, even though this new allelic form of the *DR* subregion does not code for the *DRw52* specificity. It should be noted that the DNA patterns of this novel blank specificity differ from the RFLP patterns described for another *DR* blank specificity in the family BON (Coppin et al. 1987). Analysis of family RU at the cellular and biochemical level is in progress to identify the molecules encoded by the *DR* blank genes.

(c) In family RG, the mother was typed *DR5/DR2*, whereas the father and three siblings were typed *DR5* only. This combination allowed us to compare the restriction patterns of the *DR5/5* offsprings (haplotypes *a/c* in Fig. 7) with the pattern of the father 5/- (haplotype *a/b* in Fig. 7). With the use of three enzymes, Eco RI, Bgl I, and Pst I, no difference between the banding pattern of the two *DR5/5* offspring and the banding pattern of the father was observed. These results indicate that the father does not carry any additional blank haplotype and is indeed *DR5/5* homozygous. In this example the reference to homozygous standards was not necessary.

The three examples reported for families GR, RU, and RG illustrate how the use of an analysis of *DR_β* chain genes by RFLP can allow a genetic interpretation of serological phenotypes in cases where HLA-DR typing detects only a single allele.

Acknowledgments. We thank Dr. S. Shaw for communicating unpublished results. This work was supported by the Swiss National Fund for

Scientific Research, and by CNR grant Ingeneria genetica, basi molecolari delle malattie ereditarie. C.P. was supported by EMBO and the CNRS.

References

- Albert, E. D., Baur, M. P., and Mayr, W. R.: *Histocompatibility Testing 1984*, Springer-Verlag, Berlin, 1984
- Angelini, G., de Préval, C., Ferrara, G. B., and Mach, B.: Analysis and segregation of HLA-DR by DNA typing: Correlation with supertypic specificities. In E. D. Albert, M. P. Baur, and W. R. Mayr (eds.): *Histocompatibility Testing 1984*, pp. 582-584, Springer-Verlag, Berlin, 1984
- Angelini, G., de Préval, C., Gorski, J., and Mach, B.: High-resolution analysis of the human HLA-DR polymorphism by hybridization with sequence-specific oligonucleotide probes. *Proc. Natl. Acad. Sci. U.S.A.* 83: 4489-4493, 1986a
- Angelini, G., Tanigaki, N., Tosi, R., and Ferrara, G. B.: Southern blot and microfingerprinting analysis of two *DR7* haplotypes. *Immunogenetics* 24: 63-67, 1986b
- Auffray, C. and Strominger, J. L.: Molecular genetics of the human major histocompatibility. *Adv. Immunol.* 15: 197-247, 1986
- Bell, J. I., Ester, P., St. John, T., Saiki, R., Watling, D. L., Erlich, H. A., and McDevitt, H. O.: DNA sequence and characterization of human class II major histocompatibility complex β chains from the *DR1* haplotype. *Proc. Natl. Acad. Sci. U.S.A.* 82: 3405-3409, 1985
- Böhme, J., Andersson, M., Andersson, G., Möller, E., Peterson, P. A., and Rask, L.: HLA-DR β genes vary in number between different DR specificities, whereas the number of DQ β genes is constant. *J. Immunol.* 135: 2149-2155, 1985
- Cohen, D., Le Gall, I., Marcadet, A., Font, M. P., Lalouel, J. M., and Dausset, J.: Clusters of HLA-class II β restriction fragments described allelic series. *Proc. Natl. Acad. Sci. U.S.A.* 81: 7870-7874, 1984
- Coppin, H. L., Villedieu, P., Thomsen, M., Giraud, P., Cambon-Thomsen, A., and Charron, D.: HLA-D/DR-BON: A new biochemically and cellularly defined polymorphism indistinguishable by RFLP from *DR1*. *Immunogenetics* 25: 130-132, 1987
- Gorski, J. and Mach, B.: Polymorphism of human Ia antigens: Gene conversion between two DR β loci results in a new HLA D/DR specificity. *Nature* 322: 67-70, 1986
- Gorski, J., Rollini, P., Kawashima, E., Long, E. O., and Mach, B.: Multiplicity of human class II β chain genes. In E. Sercarz, H. Cantor, and L. Chess (eds.): *Regulation of the Immune System, UCLA Symposia on Molecular and Cellular Biology, New Series*, Volume 18, pp. 47-56, Alan R. Liss, Inc., New York, 1985
- Gorski, J., Rollini, P., and Mach, B.: Structural comparison of the genes of two HLA-DR supertypic groups: The loci encoding *DRw52* and *DRw53* are not truly allelic. *Immunogenetics* 25: 397-402, 1987a
- Gorski, J., Tilanus, M., Giphart, M., and Mach, B.: Oligonucleotide genotyping shows that alleles at the *HLA-DR_βIII* locus of the *DRw52* supertypic group segregate independently of known DR or Dw specificities. *Immunogenetics* 25: 79-83, 1987b
- Grosse-Wilde, H., Doxiadis, I., and Brandt, H.: Definition of HLA-D with HTC. In E. D. Albert, M. P. Baur, and W. R. Mayr (eds.): *Histocompatibility Testing 1984*, pp. 249-264, Springer-Verlag, Berlin, 1984
- Karr, R. W.: Immunochemical analysis of the Ia polymorphisms among the family of *DR7* associated HLA-D specificities. *J. Immunol.* 136: 999-1006, 1986
- Kaufman, J., Auffray, C., Korman, A., Shackelford, D., and Strominger, J.: The class II molecules of the human and murine major histocompatibility complex. *Cell* 36: 1-13, 1984

HLA-DR

Kunkel, L. S. S.: Analysis of the human HLA-DR polymorphism by hybridization with sequence-specific oligonucleotide probes. *Proc. Natl. Acad. Sci. U.S.A.* 83: 4489-4493, 1986a

Angelini, G., Tanigaki, N., Tosi, R., and Ferrara, G. B.: Southern blot and microfingerprinting analysis of two *DR7* haplotypes. *Immunogenetics* 24: 63-67, 1986b

Auffray, C. and Strominger, J. L.: Molecular genetics of the human major histocompatibility. *Adv. Immunol.* 15: 197-247, 1986

Bell, J. I., Ester, P., St. John, T., Saiki, R., Watling, D. L., Erlich, H. A., and McDevitt, H. O.: DNA sequence and characterization of human class II major histocompatibility complex β chains from the *DR1* haplotype. *Proc. Natl. Acad. Sci. U.S.A.* 82: 3405-3409, 1985

Böhme, J., Andersson, M., Andersson, G., Möller, E., Peterson, P. A., and Rask, L.: HLA-DR β genes vary in number between different DR specificities, whereas the number of DQ β genes is constant. *J. Immunol.* 135: 2149-2155, 1985

Cohen, D., Le Gall, I., Marcadet, A., Font, M. P., Lalouel, J. M., and Dausset, J.: Clusters of HLA-class II β restriction fragments described allelic series. *Proc. Natl. Acad. Sci. U.S.A.* 81: 7870-7874, 1984

Coppin, H. L., Villedieu, P., Thomsen, M., Giraud, P., Cambon-Thomsen, A., and Charron, D.: HLA-D/DR-BON: A new biochemically and cellularly defined polymorphism indistinguishable by RFLP from *DR1*. *Immunogenetics* 25: 130-132, 1987

Gorski, J. and Mach, B.: Polymorphism of human Ia antigens: Gene conversion between two DR β loci results in a new HLA D/DR specificity. *Nature* 322: 67-70, 1986

Gorski, J., Rollini, P., Kawashima, E., Long, E. O., and Mach, B.: Multiplicity of human class II β chain genes. In E. Sercarz, H. Cantor, and L. Chess (eds.): *Regulation of the Immune System, UCLA Symposia on Molecular and Cellular Biology, New Series*, Volume 18, pp. 47-56, Alan R. Liss, Inc., New York, 1985

Gorski, J., Rollini, P., and Mach, B.: Structural comparison of the genes of two HLA-DR supertypic groups: The loci encoding *DRw52* and *DRw53* are not truly allelic. *Immunogenetics* 25: 397-402, 1987a

Gorski, J., Tilanus, M., Giphart, M., and Mach, B.: Oligonucleotide genotyping shows that alleles at the *HLA-DR_βIII* locus of the *DRw52* supertypic group segregate independently of known DR or Dw specificities. *Immunogenetics* 25: 79-83, 1987b

Grosse-Wilde, H., Doxiadis, I., and Brandt, H.: Definition of HLA-D with HTC. In E. D. Albert, M. P. Baur, and W. R. Mayr (eds.): *Histocompatibility Testing 1984*, pp. 249-264, Springer-Verlag, Berlin, 1984

Karr, R. W.: Immunochemical analysis of the Ia polymorphisms among the family of *DR7* associated HLA-D specificities. *J. Immunol.* 136: 999-1006, 1986

Kaufman, J., Auffray, C., Korman, A., Shackelford, D., and Strominger, J.: The class II molecules of the human and murine major histocompatibility complex. *Cell* 36: 1-13, 1984

- tica, basi
MBO and
- bility Test-
: Analysis
n with su-
J. R. Mayr
Springer-
- resolution
bridization
latl. Acad.
- uthern blot
types. Im-
- the human
, 1986
.. Erhlich,
acterization
hains from
1405-3409,
- son, P. A.,
n different
is constant.
- , J. M., and
fragments
J.S.A. 81:
- , Cambon-
A new bio-
inguishable
1987
gens: Gene
HLA D/DR
- Mach, B.:
az, H. Can-
tem, UCLA
es, Volume
- of the genes
DRw52 and
402, 1987a
onucleotide
ocus of the
own DR or
- of HLA-D
Mayr (eds.):
iger-Verlag,
- isms among
I. Immunol.
- i, D., and
and murine
84
- Kunkel, L. M., Smith, K. D., Boyers, H., Borgaonkar, D. S., Watchel, S. S., Millero, J., Breg, W. R., Jones, H. W., and Rary, J. M.: Analysis of human Y-chromosome-specific reiterated DNA in chromosome variants. *Proc. Natl. Acad. Sci. U.S.A.* 74: 1245-1249, 1977
- Long, E. O., Gorski, J., Rollini, P., Wake, C. T., Strubin, M., Rabourdin-Combe, C., and Mach, B.: Molecular analysis of the genes for human class II antigens of the major histocompatibility complex. *Hum. Immunol.* 8: 113-121, 1983
- Mach, B., Gorski, J., Rollini, P., Berte, C., Amaldi, I., Berdoz, J., and Ucla, C.: Polymorphism and regulation of HLA class II genes of the major histocompatibility complex. *Cold Spring Harbor Symp. Quant. Biol.* 51: 67-74, 1986
- Möller, G.: HLA and disease susceptibility. *Immunol. Rev.* 70: 1-218, 1983
- Nagy, Z. A., Baxevanis, C. N., Ishii, N., and Klein, J.: Ia antigens as restriction molecules in Ir-gene controlled T-cell proliferation. *Immunol. Rev.* 60: 59-83, 1981
- Nepom, B. S., Nepom, G. T., Mickelson, E., Antonelli, P., and Hansen, J. A.: Electrophoretic analysis of human HLA-DR antigens from HLA-DR4 homozygous cell lines: Correlation between β -chain diversity and HLA-D. *Proc. Natl. Acad. Sci. U.S.A.* 80: 6962-6966, 1983
- Nicklas, J. A., Noreen, H. J., Segall, M., and Bach, F. H.: Southern analysis of DNA polymorphism among Dw subtypes of DR4. *Hum. Immunol.* 13: 95-103, 1985
- Opelz, G.: Ninth International Histocompatibility Workshop Renal Transplant Study. In E. D. Albert, M. P. Baur, and W. R. Mayr (eds.): *Histocompatibility Testing 1984*, pp. 342-347, Springer-Verlag, Berlin, 1984
- Rollini, P., Mach, B., and Gorski, J.: Linkage map of the HLA-DR β chain genes: Evidence for a recent duplication event. *Proc. Natl. Acad. Sci. U.S.A.* 82: 7187-7201, 1985
- Schwartz, R. H.: T-lymphocytes recognition of antigens in association with gene products of the major histocompatibility complex. *Annu. Rev. Immunol.* 3: 237-261, 1985
- Se, J. K., Holbeck, S. L., Nisperos, B., Hansen, J. A., Maeda, A., and Nepom, G. T.: Identification of a polymorphic variant associated with HLA-DQw3 and characterized by specific restriction sites within the DQ β chain gene. *Proc. Natl. Acad. Sci. U.S.A.* 82: 8139-8143, 1985
- Shaw, S., Johnson, A. H., and Shearer, G. M.: Evidence for a new segregant series of B cell antigens that are encoded in the HLA-D region and that stimulate secondary allogeneic proliferative and cytotoxic responses. *J. Exp. Med.* 152: 565-580, 1980
- Southern, E. M.: Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98: 503-517, 1975
- Spielman, R. S., Lee, J., Bodmer, W. F., Bodmer, J. G., and Trowsdale, J.: Six HLA-D α -chain genes on human chromosome 6: Polymorphisms and associations of DC α -related sequences with DR types. *Proc. Natl. Acad. Sci. U.S.A.* 81: 3461-3465, 1984
- Spies, T., Sorrentino, R., Boss, J. M., Okada, K., and Strominger, J. L.: Structural organization of the DR subregion of the human major histocompatibility complex. *Proc. Natl. Acad. Sci. U.S.A.* 82: 5165-5169, 1985
- Tanigaki, N. and Tosi, R.: The genetic control of human Ia alloantigens: A three-loci model derived from the immunochemical analysis of "supertypic" specificities. *Immunol. Rev.* 66: 5-37, 1982
- Wake, C. T., Long, E. O., and Mach, B.: Allelic polymorphism and complexity of the genes for HLA-DR β -chains-direct analysis by DNA-DNA hybridization. *Nature* 300: 372-374, 1982

Received March 16, 1987